

2016 ACROSS-BREED EPD ADJUSTMENTS

The table below contains adjustment factors to compare EPDs of **individuals** of different breeds, based on studies by the U. S. Meat Animal Research Center, Clay Center, Nebraska. Adjustments for some breeds are not available if they are not in the MARC database. The base for comparison is Angus.

These factors should be added to an individual EPD for direct comparison with an individual EPD of a different breed. For instance, consider an Angus sire with Weaning Weight EPD of +50 and a Charolais sire with Weaning Weight EPD of +25. The Charolais adjustment factor is +35, so the adjusted value for the Charolais sire is +60 (25 + 35), 10 lb heavier than the Angus sire's +50 (which is not adjusted since Angus is the basis of comparison). Or, compare a Gelbvieh sire with Milk EPD of +23 with a Hereford sire of +20. The Gelbvieh adjustment factor is +2 and the Hereford adjustment factor is -17; the result is the Gelbvieh sire's adjusted value is +25 (23 + 2) and the Hereford adjusted value is +3 (20 - 17), 23 lb below the Gelbvieh.

These factors are not breed comparisons. They are adjustments to allow comparison of **individuals** of different breeds. Breed comparisons can be seen on this website (<http://beef.tamu.edu>) in the list of publications under "Genetics & Selection". Look for "2016 Sire-Breed Comparisons for EPD Traits". Registry association breed averages can also be seen in another publication on the above website under "2016 Breed Averages for EPD Traits".

Other EPD values calculated by some breeds (but for which there are no across-breed adjustment factors) include various combinations of direct and maternal calving ease, gestation length, scrotal circumference, yearling hip height, mature weight, mature height, docility, stayability (longevity in the herd), Yield Grade or cutability (estimate percent lean), tenderness, and \$ Value Indexes. For a complete breed genetic evaluation report, contact the particular breed association or access their website.

| Breed | Birth Wt | Wean Wt | Yearling Wt | Milk | Marbling | REA | Fat Cover |
|------------|----------|---------|-------------|------|----------|-------|-----------|
| Angus | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Beefmaster | +4.7 | +18 | 1 | +6 | | | |
| Brahman | +10.3 | +45 | +7 | +24 | -0.78 | -0.05 | -0.14 |
| Brangus | +3.3 | +12 | +4 | +6 | | | |
| Braunvieh | +1.9 | -26 | -50 | -1 | -0.70 | +0.78 | -0.09 |
| Charolais | +8.0 | +35 | +40 | +8 | -0.33 | +1.01 | -0.21 |
| Chiangus | +3.2 | -27 | -40 | -2 | -0.34 | +0.34 | -0.09 |
| Gelbvieh | +2.8 | -23 | -29 | +2 | -0.27 | +0.75 | |
| Hereford | +2.3 | -8 | -29 | -17 | -0.31 | -0.07 | -0.06 |

| | | | | | | | |
|-----------------|------|-----|-----|-----|-------|-------|-------|
| Limousin | +2.3 | -18 | -41 | -14 | -0.43 | +1.01 | -0.13 |
| MaineAnjou | +4.2 | -30 | -39 | -6 | -0.57 | +0.97 | -0.19 |
| Red Angus | +2.5 | -31 | -35 | +3 | -0.27 | +0.01 | -0.02 |
| Salers | +1.8 | -7 | -26 | +6 | -0.09 | +0.94 | -0.20 |
| Santa Gertrudis | +5.7 | +36 | +43 | +17 | -0.54 | -0.09 | -0.08 |
| Shorthorn | +4.7 | -37 | -17 | +4 | -0.14 | +0.37 | -0.10 |
| Simmental | +3.2 | -9 | -12 | +4 | -0.34 | +0.49 | -0.13 |
| South Devon | +3.3 | -11 | -27 | +4 | -0.08 | +0.29 | -0.13 |
| Tarentaise | +3.4 | +25 | +6 | +24 | | | |

Updated October, 2016